



University Animal Care Committee Standard Operating Procedure		
Document No: 10.6	Subject: Anesthesia in Rats	
Date Issued: March 14, 2012	Revision: 2	Page No: 1

Location: Queen's University

Responsibility: Principal Investigators (PI), Research Staff, Veterinary Staff

Purpose: The purpose of this Standard Operating Procedure (SOP) is to describe approved methods commonly used to anesthetize rats.

1. Introduction and Definitions: The anaesthetic regimen must be compatible with the needs and restrictions of the procedure to be performed, taking into account the length of the procedure, whether or not the animal is expected to survive the procedure, the degree of invasiveness and the scientific goals.

Fasting prior to anaesthesia is generally not required in rats as vomiting is extremely rare during induction and anaesthesia. If scientifically justified, fasting in rats should be as short as possible due to the high metabolic rate of these small mammals.

Either inhalant or injectable anaesthetics can be used to anaesthetise rats. Inhalant anaesthetics with the use of a vaporiser are the agents of choice as they provide a safe, reliable, reversible, and reproducible method.

Once anaesthetized, ophthalmic ointment must be instilled to prevent drying of the eye in the (e.g. Hypotears, Lubrithal Eye Gel).

Monitor the rat continuously as an anaesthetized animal should never be left unattended.

As rodents have a high surface area to body mass ratio, it is extremely important to provide thermal support starting immediately after induction of anaesthesia and continuing until full recovery.

Fluid therapy must be provided after each and every anesthetic event (regardless of the procedure performed). The only exception is if fluid may confound results; this deviation must then be described within the approved Animal Use Protocol.

Abbreviations: intravenous **IV**, subcutaneous **SC**, intraperitoneal **IP**, intramuscular **IM**, per os **PO**

2. Materials:

- Warm water recirculating blanket, heating disks etc.
 - Gas anaesthesia machine (calibrated within the last 12 months, and with scavenging capabilities or a passive filter)
 - Transparent induction chamber
 - Nosecone
 - Isoflurane
 - Ketamine 100mg/kg (controlled substance – exemption permit required)
-



University Animal Care Committee Standard Operating Procedure

Document No:
10.6

Subject:
Anesthesia in Rats

Date Issued:
March 14, 2012

Revision:
2

Page No:
2

- Xylazine (20 mg/ml)
 - Acepromazine (10mg/ml)
 - Medetomidine
 - Atipamezole
 - Sterile sodium chloride 0.9% or sterile water (diluent)
 - Sterile sodium chloride 0.9% or Lactated Ringers solution (fluid therapy)
 - Sterile needles (25-27gauge)
 - Sterile syringes (various sizes)
 - Alcohol swabs
 - Bell jar (or other suitable transparent container with lid)
 - Gauze
 - Eye lubricant
 - 10ml red top vacutainers

 - **Procedures for Adult Rats:**
 - ***Isoflurane Anaesthesia:***
 - i. Induction
 1. Place the animal in the induction chamber.
 2. Adjust the oxygen flow to 1 – 1.5 L/min.
 3. Adjust the vaporiser to 3% - 5%.
 4. Once the rat is in lateral recumbency and cannot right itself, apply eye lubricant and transfer the animal to a nosecone for maintenance.
 - ii. Maintenance
 1. Ensure the rat's snout is firmly seated in the nosecone with a good seal.
 2. Adjust the oxygen flow to ~1L/min.
 3. Adjust the vaporiser to 1 – 2.5% (this is variable depending on the depth of anesthesia required for the particular procedure).
 4. Maintain the rat on a heating device for the duration of the procedure.
 - iii. Recovery
 1. Turn off the isoflurane and leave the animal on 100% oxygen (short term).
 2. Administer fluid therapy.
 3. Transfer the animal to a clean cage and place it on a piece of paper towel until it is mobile.
 4. Provide heat during this recovery phase.
 5. Return the animal to the colony room once it has fully recovered from anaesthesia (blinking and righting reflex have returned).

 - ***Ketamine/Xylazine Anaesthesia:***
 - i. Anaesthetic dose: ketamine 75 mg/kg, xylazine 10 mg/kg
 - ii. To prepare the cocktail, in a sterile vial mix:
 1. 0.75ml (75 mg) ketamine (100 mg/ml)
-



University Animal Care Committee Standard Operating Procedure		
Document No: 10.6	Subject: Anesthesia in Rats	
Date Issued: March 14, 2012	Revision: 2	Page No: 3

2. 0.25 ml (5 mg) xylazine (20 mg/ml)
 3. 0.75 ml diluent
 4. Label with the drug name(s), expiration date of the drugs, date mixed and initials.
- iii. Gently restrain the animal, swab the injection site with 70% alcohol and administer 0.2ml per 100g of body weight IP.
 - iv. Duration of anaesthesia is 20 – 40 minutes depending on the strain, sex, and body weight of the animal.
 - v. If a top up dose is required, a half dose of only ketamine (38 mg/kg) should be administered. To prepare the half dose:
 1. 0.38 ml (37.5 mg) ketamine (100 mg/ml)
 2. 4.62 ml diluent
 3. Write on the vial the name of the drug, expiration date, date mixed and initials
 - vi. Gently restrain the animal, swab the injection site with 70% alcohol and administer 0.2ml per 100g body weight IP.
 - vii. Atipamezole (5 mg/ml) can be administered to reverse xylazine and facilitate recovery at a dose of 0.1 to 5.0 mg/kg SC or IP.
- ***Ketamine/Xylazine/Acepromazine Anaesthesia:***
 - i. Anaesthetic dose: ketamine 50 mg/kg, xylazine 5 mg/kg, acepromazine 1 mg/kg.
 - ii. To prepare cocktail, in a sterile vial or bottle with a rubber stopper, mix:
 1. 5 ml (500 mg) ketamine (100 mg/ml)
 2. 2.5 ml (50 mg) xylazine (20 mg/ml)
 3. 1 ml (100 mg) acepromazine (10 mg/ml)
 4. 1.5 ml diluent
 5. Label with the drug name(s), expiration date of the drugs, date mixed and initials.
 - iii. Mixed cocktail should be protected from light and stored in a cool place.
 - iv. Gently restrain the animal, swab the injection site with 70% alcohol and administer 0.1ml per 100g body weight IP.
 - v. Duration of anaesthesia is approximately 30 minutes depending on the strain, sex, and body weight of the animal.
 - vi. After 30 minutes, a half dose (0.05 ml per 100g) may be administered as needed.
 - ***Ketamine/Medetomidine Anaesthesia:***
 - i. Anaesthetic dose: ketamine 75 mg/kg, medetomidine 0.5 mg/kg
To prepare cocktail, in a sterile vial or bottle with a rubber stopper, mix:
 1. 3 ml (300 mg) ketamine (100 mg/ml)
 2. 2 ml (2 mg) medetomidine (1mg/ml)
 3. 1 ml diluent
 4. Label with the drug name(s), expiration date of the drugs, date mixed and initials
-



University Animal Care Committee Standard Operating Procedure		
Document No: 10.6	Subject: Anesthesia in Rats	
Date Issued: March 14, 2012	Revision: 2	Page No: 4

- ii. Mixed cocktail should be protected from light and stored in a cool place.
- iii. Gently restrain the animal, swab the injection site with 70% alcohol and administer 0.15 ml per 100g body weight IP.
- iv. Duration of anaesthesia is approximately 30 minutes depending on the strain, sex, and body weight of the animal.
- v. If a top up dose is required, only ketamine (75 mg/kg) should be administered. To prepare the dose:
 1. 0.38 ml (37.5 mg) ketamine (100 mg/ml)
 2. 4.62 ml diluent
 3. Label with the drug name, expiration date, date mixed and initials
- vi. Gently restrain the animal, swab the injection site with 70% alcohol and administer 0.2ml per 100g body weight IP.
- vii. Atipamezole (5 mg/ml) can be administered to reverse xylazine and facilitate recovery at a dose of 0.1 to 5.0 mg/kg SC or IP.

- ***Bell Jar Anaesthesia:***

- i. As this method of anesthesia has a high mortality risk, bell jar anesthesia must be referenced and explicitly described in the active Animal Use Protocol (Anesthetic Agent table and Euthanasia Methods). The following must be implemented:
 1. The bell jar must be used in a hood.
 2. The isoflurane must be poured onto clean gauze, which is placed within the bell jar.
 3. The grid of the bell jar, or another device (such as an upside-down petri dish) must be placed on the soaked gauze so the animal has no direct contact with the volatile compound.
 4. Only one animal at a time is permitted in the bell jar, and it must have adequate room.
 5. The bell jar must be covered so the animal cannot escape during the excitatory phase.
 6. The bell jar must be kept clean.

Revised: February 28, 2019 / June 20, 2019
